

E17-2004-175

D. I. Uzunov\*

A NEW TREATMENT  
OF FLUCTUATION CORRELATIONS  
NEAR PHASE TRANSITION POINTS

---

\*G. Nadjakov Institute of Solid State Physics, BAS, BG-1784 Sofia,  
Bulgaria; [uzun@issp.bas.bg](mailto:uzun@issp.bas.bg)

Узунов Д. И.

E17-2004-175

Новое рассмотрение флуктуационных корреляций вблизи точки фазового перехода

Рассмотрен общий самосогласованный подход, с помощью которого возможно полное рассмотрение поправок к стандартному среднеполевому приближению. Естественное обобщение стандартного приближения среднего поля с помощью разложения кумулянтного вида приводит к новому пониманию теории среднего поля. Предложенный подход может быть применен для систематического рассмотрения флуктуационных эффектов разного масштаба длин и для разработки (развития) нового метода статистического огрубления. Получены конкретные результаты для критической температуры и параметров Ландау в  $\phi^4$ -теории, представляющей полевой эквивалент модели Изинга. В рамках предложенного подхода рассмотрено вычисление истинной критической температуры — вопрос, который является нерешенным в современной теории фазовых переходов.

Работа выполнена в Лаборатории теоретической физики им. Н. Н. Боголюбова ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна, 2004

Uzunov D. I.

E17-2004-175

A New Treatment of Fluctuation Correlations Near Phase Transition Points

A general self-consistency approach which allows a thorough treatment of the corrections to the standard mean-field approximation (MFA) has been reviewed. The natural extension of standard MFA with the help of a cumulant expansion leads to a new point of view on the MF theories. The proposed approach can be used for a systematic treatment of fluctuation effects of various length scales and, perhaps, for the development of a new coarse-graining procedure. We outline and justify our method by some preliminary calculations. Concrete results are given for the critical temperature and the Landau parameters of the  $\phi^4$ -theory – the field counterpart of the Ising model. An important unresolved problem of the modern theory of phase transitions – the problem for the calculation of the true critical temperature, is considered within the framework of the present approach. A comprehensive description of the ground state properties of many-body systems is also demonstrated.

The investigation has been performed at the Bogoliubov Laboratory of Theoretical Physics, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna, 2004

## INTRODUCTION

The main aim of this paper is focused on the correspondence between microscopic models of phase transitions and their quasi-macroscopic (field-theoretic) counterparts. Here we shall outline a new self-consistency approach to a more accurate derivation of effective field theories from microscopic models defined on lattices.

Our method is general and can be used for a wide class of microscopic models but for a concreteness here we shall illustrate our approach with the Ising model (IM), given by

$$\mathcal{H}(s) = -\frac{1}{2} \sum_{ij}^N J(|i-j|) s_i s_j, \quad (1)$$

where  $s \equiv \{s_i\}$  denotes a lattice «field»,  $s_i = \pm 1$ , and the interaction constant of ferromagnetic type  $J_{ij} = J(|i-j|) > 0$  depends on the intersite distance  $|i-j|$  in a regular  $D$ -dimensional lattice of  $N$  sites («spins» or pseudo-spins). Note, that  $J(0) \equiv J_{ii} = 0$ .

We shall follow the main path of the phase transition theory, where the effective (quasi-macroscopic) field Hamiltonians (alias Ginzburg–Landau (GL) free energies) are derived with the help of two systematic methods: i) Hubbard–Stratonovich transformations (HST) and ii) a mean-field (MF) like procedure [1–3]. Here we propose a more thorough approach based on a convenient generalization of (ii).

The known field theories exhibit both success and failure in the description of many-body systems defined by microscopic models. For example, we believe that the renormalization group methods of the modern theory of phase transitions [1] based on the  $\phi^4$ -theory yield a quite convenient description of the scaling and universality properties of IM but we cannot be certain that the field theory satisfactorily describes important non-universal properties, such as, for example, the critical temperature  $T_c$  and the lower critical dimensionality  $D_L$  (for IM,  $D_L = 1$ , whereas within the  $\phi^4$ -theory,  $D_L = 2$ ). In fact the field theory fails along this line of studies. For example, the fluctuation shift  $(\Delta T_c)_f$  of  $T_c$  predicted within the one-loop approximation for the  $\phi^4$ -theory is a very small and, hence, unrealistic, while in the higher orders of the loop expansion this shift turns out infinite and its calculation needs a special renormalization. As a result,

the problem for the value of  $T_c$  within the framework of the present field theory of phase transitions remains unresolved. Here we shall show the genesis of this problem and present a satisfactory solution.

The reason for the mentioned difficulties of the field theory approach is in the quite simplified coarse-graining procedures used in the derivation of the GL effective Hamiltonians from microscopic models. HST is applied together with the long-wavelength approximation (LWLA),  $(ka_0) \ll \pi$ ;  $k = |\mathbf{k}|$ ;  $\mathbf{k} = (k_1, \dots, k_D)$  is the wave number,  $a_0$  is either the lattice constant or, generally, the mean interparticle distance. LWLA leads to a correct expression for the Ornstein–Zernicke correlation function but cuts the short-range (i.e. high-energy:  $\epsilon(k) \sim k^2, \Lambda < k < \phi/a_0$ ) interparticle correlations of fluctuation type that have the main contribution to the shift  $(\Delta T_c)_f$ ;  $\Lambda \ll \pi/a_0$  is the upper cutoff for  $k$  within LWLA.

In the approach (ii) we usually say that we neglect the fluctuations of the physical quantities from their equilibrium values. This is not entirely true. It seems important to emphasize that the latter are values in MF approximation (MFA) and, hence, they are incorrect. Thus the fluctuations are defined towards incorrect statistical averages, and their contribution to the free energy of the system cannot be accepted as an entire fluctuation effect. The latter can be correctly evaluated, if we are able to define the fluctuations as variations towards exact statistical averages calculated by Hamiltonian (1). This task seems unsolvable, but the present paper makes a step of improvement of the theory along the same direction.

Note, that the methods (i) and (ii), in spite of the difference between them, lead to the same GL effective field theory. Both methods use LWLA. For the method (ii) LWLA seems to me obligatory because of the following important argument. HST can be applied only to positively definite matrices  $(J_{ij})$  but this condition is not satisfied by quite important lattice models, such as the nearest-neighbour ( $nn$ ) IM. In LWLA, however, the same interaction matrix  $J_{ij;nn}$  of IM is modified to a form that is positively definite. Besides, in both methods, LWLA is used to help the derivation of the quasi-macroscopic (fluctuation) and macroscopic (thermodynamic) properties, i.e., the LWLA is a tool of a «coarse-graining» procedure for the many-body system. But in both cases (i) and (ii), the micro- and mesoscopic interparticle correlations are ignored. Here we shall use a generalization of the method (ii) in order to improve this disadvantage of the theory. In this way we shall present more accurate calculation of the LG parameters (vertices) of the effective field theory of Ising systems. Besides we shall show explicitly for the first time the mechanism of «statistical correlation» which leads from the two-site ( $i - j$ ) trivial correlation presented by the initial interaction  $J_{ij}$  to short-, meso-, and large-scale effective multiparticle correlations of fluctuation type — fluctuation correlations. Thus we shall establish and develop for the first time a new «coarse-graining» procedure, and this is the main aim of

our work. Besides, we shall show that the GL parameters of the  $\phi^4$ -theory acquire  $(1/z)$ -corrections for both short-range and long-range interactions  $J(|i - j|)$  [1]. These corrections will be presented to the second order in  $(1/z)^2$ .

The  $1/z$ -expansion has been introduced by R. Brout [4] and applied in calculations of  $T_c$  and thermodynamic susceptibilities [5–13]. We are not aware of another relevant work along this line of research except for our recent investigation [2, 3]; the latter will be used in our investigation. Remember, that the (Brout) approach is a development of an older method – the Kirkwood method of cumulants (semi-invariants); hence, we shall follow a cumulant expansion which is well known.

We shall generalize the Brout approach in a way that makes it possible to derive effective field theories, and this will allow us to reveal new and surprising features of many-body systems. In the Brout scheme, the mean («molecular») field is spatially ( $i$ -) independent. We find that there are no physical reasons for this assumption and extend the «mean-field concept». Within our approach, the so-called mean field is spatially dependent up to the moment when one should find the actual ground state. Then the fluctuation phenomena occur with respect to this ground state. In our approach the GL parameters and, hence, the ground state have a more precise evaluation.

Our consideration has been performed for interactions  $J(|i - j|)$  of a quite general type, namely for all interactions that can be presented by the equality

$$\sum_j J_{ij} = zJ_0 \equiv J, \quad (2)$$

where  $J_0$  is an effective exchange constant, and  $z$  is an effective «coordination» number (number of interacting neighbours). In particular cases our results will be referred to the most common case of  $nn$  interactions (then  $n = 2D$  for simple lattices). For a simplicity, here we shall assume that IM is defined by a simple cubic (*sc*) lattice. The consideration can be easily expended to other types of regular lattices, as well as to irregular lattices with certain forms of quenched disorder, for example, random potential [1]. We introduce the interaction radius by  $R_{\text{int}} \approx z^{1/D}$  – a quantity, which is equal to the so-called zero-temperature correlation length (see Sec. 1). Let us emphasize that our results can be rederived without difficulties in the presence of an external field  $\{h_i\}$  conjugate to the lattice field  $\{s_i\}$ .

In Sec. 1 we present a general approach to the treatment of fluctuation correlations at various length scales and discuss aspects of the usual theory that corresponds to the lowest order approximation of a perturbation expansion of cumulant type. In Sec. 2 we investigate the higher orders of the mentioned perturbation expansion and demonstrate several new features of the effective field theory. Our main results are summarized and discussed in Subsecs. 2.4–2.7. This work will be published in a more extended form [15].

## 1. PRESENT STATUS OF THE EFFECTIVE FIELD THEORY

**1.1. General Scheme.** The equilibrium free energy of IM as a function of the temperature  $T$  and the field configuration  $h = \{h_i\}$  is given by

$$G(T) = -\beta^{-1} \ln \left\{ \text{Tr} e^{-\beta \mathcal{H}(s)} \right\}, \quad (3)$$

where  $\beta^{-1} = k_B T$ , and the Trace is over the allowed lattice configurations  $s = \{s_i\}$ . Note, that the equilibrium values of the physical quantities are calculated as averages with respect to the statistical ensemble based on Hamiltonian (1) and, in particular, the statistical averages of type  $\langle s_i \dots s_j \rangle$  are obtained as derivatives of the partition sum whereas the irreducible averages  $\langle\langle s_i \dots s_j \rangle\rangle = \langle (s_i - \langle s_i \rangle) \dots (s_j - \langle s_j \rangle) \rangle$  are obtained as derivatives of the Gibbs free energy (3).

Let us introduce the shift

$$s_i = \phi_i + \delta s_i, \quad (4)$$

where the lattice field  $\phi_i$  is an arbitrary (auxiliary) field configuration that is not necessarily associated with the averaged spin  $\langle s_i \rangle$  at site  $i$ , and the «fluctuation»  $\delta s_i$  is merely the difference  $(s_i - \phi_i)$ . The identification of  $\phi_i$  with a statistical average  $\langle \dots \rangle$  over the full Hamiltonian (1), or, with a statistical average  $\langle \dots \rangle_0$  corresponding to another ensemble as well as the interpretation of  $\delta s_i$  as a fluctuation around  $\langle s_i \rangle$  (or  $\langle s_i \rangle_0$ ) may be a matter of further considerations. At this stage  $\phi_i$  and  $\delta s_i$  are auxiliary variables which obey (4) and are not referred to concrete physical quantities.

Following a standard procedure (see, e.g., Refs. [1, 2]) we obtain the following effective non-equilibrium free energy:

$$\mathcal{H}(\phi) = \mathcal{H}_0(\phi) + \mathcal{H}_f(\phi), \quad (5)$$

with  $\phi \equiv \{\phi_i\}$ ,

$$\mathcal{H}_0(\phi) = \frac{1}{2} \sum_{ij} J_{ij} \phi_i \phi_j - \beta^{-1} \sum_i \ln [2 \text{ch}(\beta a_i)]. \quad (6)$$

Here

$$a_i = \sum_j J_{ij} \phi_j \quad (7)$$

is the «mean» (molecular) field, and

$$\mathcal{H}_f(\phi) = -\beta^{-1} \ln \left\langle \exp \left[ \frac{\beta}{2} \sum_{ij} J_{ij} (s_i - \phi_i)(s_j - \phi_j) \right] \right\rangle_0 \quad (8)$$

is the «fluctuation» part. As usual, we shall often call the free energy (5) an «effective Hamiltonian».

In (8),  $\langle \dots \rangle_0$  denotes a statistical average over an ensemble defined by the auxiliary («MF») Hamiltonian

$$\mathcal{H}_a(\phi, s) = - \sum_i a_i s_i; \quad (9)$$

the respective partition function and (nonequilibrium) free energy are given by  $\mathcal{Z}_a(\phi) = \text{Tr}[\exp(-\beta\mathcal{H}_a)]$ , and  $G_a(\phi) = -\beta^{-1} \ln \mathcal{Z}_a(\phi)$ . For this simple ensemble, we have

$$\langle s_i \rangle_0 = \text{th} [\beta a_i(\phi)]. \quad (10)$$

The calculation of averages of type  $\langle s_i \dots s_j \rangle_0$  as well as «irreducible» averages of type  $\langle \langle \delta s_i \dots \delta s_j \rangle \rangle_0$  is also straightforward. These averages represent a form of fluctuation correlations, but they are just an auxiliary theoretical tool rather than real objects. In contrast, the real objects, namely, (full) statistical averages  $\langle \dots \rangle$  and  $\langle \langle \dots \rangle \rangle$  within the total Hamiltonian (1) cannot be exactly calculated.

**1.2. Usual Theory.** In the framework of the usual theory, the «fluctuation» term  $\mathcal{H}_f$  is ignored. This is the MFA. In the present general format of the theory, we have  $N$  (self-consistency) equations of state  $(\partial\mathcal{H}/\partial\phi_i) = 0$  at fixed  $T$  (and  $h = \{h_i\} = 0$ ) — one equation per a lattice vertex  $i$ :

$$\sum_j J_{ij} \{ \bar{\phi}_j - \text{th} [\beta a_j(\bar{\phi})] \} = 0, \quad (11)$$

where  $\bar{\phi} = \{\phi_i\}$ . It is easy to see that the number  $(z-1)$  of nonzero terms ( $i \neq j$ ) in all  $N$  sums (11) is equal to the number of nonzero interaction constants acting on the site  $i$ :  $J(|i-j|) > 0$  for  $a_0 \leq |i-j| \leq R_{\text{int}}$ ;  $J(|i-j|) = 0$  for  $|i-j| \equiv R > R_{\text{int}}$ . The «equations of state» (11) can be written in the simple form

$$\bar{\phi}_i = \text{th} [\beta a_i(\bar{\phi})]. \quad (12)$$

The equivalence of (11) and (12) can be easily proven for any number  $N \geq 1$ .

From (10) and (12) we obtain

$$\langle \bar{s}_i \rangle_0 = \text{th} [\beta a_i(\bar{\phi})] = \bar{\phi}_i. \quad (13)$$

Thus in this quite general form of MFA ( $\mathcal{H}_f \approx 0$ ), we have:  $\langle s_i \rangle_0 = \bar{\phi}_i$ ,  $\langle \dots \rangle = \langle \dots \rangle_0$ , and  $G(T, h) = \mathcal{H}_0(\bar{\phi})$  is the equilibrium free energy that corresponds to extrema (including minima)  $\bar{\phi}$  of  $\mathcal{H}(\phi) \approx \mathcal{H}_0(\phi)$  — the non-equilibrium MF free energy given by the non-equilibrium (arbitrary) order parameter field  $\phi_i$ . By  $\bar{\phi}_i$  from (11) we denote the equilibrium configuration of the latter; hereafter the «bar» of the equilibrium value  $\bar{\phi}_i$  of  $\phi_i$  will be often omitted. Now one may

perform a Landau expansion for small  $\bar{\phi}_i$  in order to get other known forms of the MF theory.

However, at the present stage of consideration we are interested in some more generality and for this reason we continue our discussion of the non-equilibrium free energy functional

$$\tilde{G}(T/\phi) \equiv \mathcal{H}(T/\phi) \approx \mathcal{H}_0(T/\phi) \quad (14)$$

as given by Eq. (6). In (14), the dependence of the non-equilibrium free energy  $\tilde{G}$  on  $\phi$  is denoted by «/ $\phi$ » because of the more special role of this variable, namely, the variation  $\delta\tilde{G}$  should be zero at thermal equilibrium and from this condition one obtains the possible thermal equilibria  $\bar{\phi}$  (alias «self-consistency condition»).

The expansion of the *log*-term in (6) up to fourth order in  $\phi_i$  yields the known result for the lattice  $\phi_i^4$ -theory of the Ising model:

$$\begin{aligned} \mathcal{H}_0(\phi) = & \frac{1}{2} \sum_{ij} J_{ij} \phi_i \phi_j - \frac{\beta}{2} \sum_{ijk} J_{ij} J_{ik} \phi_j \phi_k + \\ & + \frac{\beta^3}{12} \sum_{ijklm} J_{ij} J_{ik} J_{il} J_{im} \phi_j \phi_k \phi_l \phi_m. \end{aligned} \quad (15)$$

One may apply LWLA to this form of the theory but we shall follow a different path.

**1.3. Continuum Limit.** Using the rule

$$\sum f_i = \rho \int d^D x f(\mathbf{x}) \equiv \rho \int d\mathbf{x} f(\mathbf{x}), \quad (16)$$

where  $\rho = (N/V)$  we can write Eq. (2) in the form

$$J = \rho \int d^D R J(R), \quad (17)$$

with  $R = |\mathbf{R}|$ ;  $\mathbf{R} = (\mathbf{x} - \mathbf{y})$ . Note, that the factor  $\rho$  in (19)–(21) can be avoided and this is the usual practice. In the latter case the physical dimension of the respective physical quantity is changed by a factor  $[V] \sim [L]^D$ ; for example,  $[L]^D [f_i] = [f(\mathbf{x})]$ . Of course, one may use both variants.

In the continuum limit effective Hamiltonian (6) corresponding to a zero external field ( $h = 0$ ) takes the form

$$\mathcal{H}_0 = \frac{\rho}{2} \int d^D x \phi(\mathbf{x}) I[\phi(\mathbf{x})] - \rho \beta^{-1} \int d^D x \ln \text{ch} \{ \beta^{-1} I[\phi(\mathbf{x})] \}, \quad (18)$$



where

$$I[\phi(\mathbf{x})] = \rho \int d^D y J(R) \phi(\mathbf{y}). \quad (19)$$

Now we apply LWLA in the form

$$[\phi(\mathbf{x}) - \phi(\mathbf{y})]^2 \ll |\phi(\mathbf{x})\phi(\mathbf{y})|, \quad (20)$$

and under this assumption truncate the Taylor expansion

$$\phi(\mathbf{y}) = \phi(\mathbf{x}) + \sum_{\alpha=1}^D \frac{\partial \phi(\mathbf{x})}{\partial x_\alpha} R_\alpha + \frac{1}{2} \sum_{\alpha, \beta=1}^D \frac{\partial^2 \phi(\mathbf{x})}{\partial x_\alpha \partial x_\beta} R_\alpha R_\beta + \dots \quad (21)$$

to the second order in  $\mathbf{R} = \{R_\alpha\}$ .

Using approximation (20), (19) becomes

$$I[\phi(\mathbf{x})] = J\phi(\mathbf{x}) + \frac{\tilde{J}}{2D} \nabla^2 \phi(\mathbf{x}), \quad (22)$$

where  $J$  is given by (17), and

$$\tilde{J} = \rho \int d^D R J(R) R^2. \quad (23)$$

Now we substitute (22) in (18), perform the expansion up to order  $\phi^4(\mathbf{x})$  and to the second order in  $\nabla \phi(\mathbf{x})$ . Besides, we should keep in mind, that in expansion in powers of  $\phi_i$  we cannot distinguish between  $T$  and  $T_{c_0}$  except for the  $\phi_i^2$ -term where the difference between  $T$  and  $T_{c_0}$  should be kept only to the lowest nonvanishing order; in our case, this is the first order in  $(T - T_{c_0})$  (see, e.g., Ref. [1]). Following these notes, we perform at a certain stage of the calculation an integration by parts with the convenient boundary condition  $\nabla \phi(\mathbf{x}) = 0$  and obtain the well-known GL effective Hamiltonian

$$\mathcal{H}_0[\phi(\mathbf{x})] = \rho \int d^D x \left\{ \frac{\tilde{c}_0}{2} [\nabla \phi(\mathbf{x})]^2 + \frac{r_0}{2} \phi^2(\mathbf{x}) + u_0 \phi^4(\mathbf{x}) \right\}, \quad (24)$$

with

$$c_0 = \frac{R_{\text{int}}^2}{2D} J, \quad r_0(T) = k_B(T - T_{c_0}), \quad u_0 = \frac{J}{12}. \quad (25)$$

Here  $T_{c_0} = (J/k_B)$  and terms of order  $t_0 = (T - T_{c_0})/T_{c_0} \ll 1$  have been neglected in  $c_0$  and  $u_0$  [2], i.e. these two parameters are calculated at  $T_{c_0}$  [1, 2].

To clarify the result (24)–(25) we shall mention that  $J(R)$  for  $R > R_{\text{int}}$  is very small and can be ignored. Setting  $J(R) \sim J_0$  in (17), comparing the result with  $J = zJ_0$  from (2), and noticing that  $\rho = 1/v \sim a_0^D$ , one obtains

$z \sim (R_{\text{int}}/a_0)^D$  as should be, and  $J \approx J_0 R_{\text{int}}^D$ . In the same way one gets  $\tilde{J} \approx J R_{\text{int}}^2$ .

The results (24)–(25) show that the energy of the spatially dependent configurations of the field depends on the interaction radius. The latter serves as a coherence (correlation) length of the field  $\phi(\mathbf{x})$ ; see the parameter  $c_0$  given in (25). In order to clarify this point, let us consider the so-called zero-temperature correlation length [1], defined by  $\xi_0 \equiv \xi(T=0) = [-c_0/r_0(0)]^{1/2}$ . Using (25) and  $T_{c_0} = J/k_B$  we obtain  $\xi_0 = R_{\text{int}}\sqrt{2D}$ .

**1.4. Discussion.** Note that the temperature range of validity of these considerations is  $t_0(T) \ll 1$  and  $(ka_0) \ll \pi$ . The spatially dependent fluctuations correspond to a higher energy than the uniform configuration, and hence the latter contains the deepest (global) minima  $\bar{\phi}$  of the effective free energy, nevertheless we have written  $N$  «equations of state» as a result of the minimization of the effective Hamiltonian. Now one can easily show that the variation of Hamiltonian (24) with respect to the field will give again spatially dependent solutions but in the usual theory they are interpreted as «spatially dependent fluctuations» which, together with the uniform fluctuation  $\delta\phi = (\phi - \bar{\phi})$  towards the stable state  $\bar{\phi}$ , are all fluctuations in the system in LWLA. But we know that this picture contains the approximation  $\langle s_i \rangle \sim \langle s_i \rangle_0 = \bar{\phi}$ .

Our point of view is the following. The averages  $\langle \rangle_0$  should not be taken very seriously. They are an auxiliary tool in our consideration, and are not the final aim of our investigation. We have used these averages only because they appear along our way of obtaining an effective Hamiltonian  $\mathcal{H}$  (or  $\mathcal{H}_0$  in the lowest order of the theory) in which the statistical degree of freedom  $\phi_i$  varies in a wide range of values ( $-\infty < \phi_i < \infty$ ). This is the only consistent interpretation of our consideration performed so far. We can assume that up to now we have obtained nothing else but an effective free energy (effective lattice Hamiltonian) in terms of the new lattice field  $\phi_i$ . In a clear approximation, this effective model is given by (6); for the expansion in powers of  $\phi_i$ , see (15).

Another important aspect of our consideration is that the new Hamiltonian contains more  $ij$ -interactions than original model (1). The mathematical form of (15) gives a clear physical interpretation of these interactions: the first  $\phi_i\phi_j$ -term in the r.h.s. of (15) describes an interaction that is quite similar to the original inter-spin interaction in (1) whereas the second term of the same type in (15) describes an indirect two-site  $jk$ -interaction that is mediated by the « $i$ -spins». This means that the latter interaction  $J_{ij}J_{ik}$  has twice larger radius of action than the original  $J_{ij}$  exchange. The four-point interaction given by the third term in the r.h.s. of (15) can also be described in the above style, introduced for the first time in this paper.

Therefore, even at this early stage of consideration ( $\mathcal{H} \approx \mathcal{H}_0$ ) we see that the effective free energy exhibits effects of «statistical extension of the original

inter-particle correlations (interactions)». This «principle of growth of statistical correlations» is well known in the general phase transition theory. Here we show the concrete mechanism of the respective phenomenon and a systematic way of description of growth of statistical correlations. This point will become more clear from the results in the next Section. We shall see that the investigation along this path leads to a quite unexpected and intriguing picture.

## 2. BEYOND THE STANDARD THEORY

**2.1. Perturbation Series.** Here we consider the  $\phi$ -contributions to the effective free energy (Hamiltonian)  $\mathcal{H}(\phi)$  which are generated by the term  $\mathcal{H}_f(\phi)$ . For obvious reasons, terms that are  $\phi_i$ -independent will be omitted.

It seems convenient to rearrange our theory by introducing the auxiliary variables  $\Delta_i = (s_{i0} - \phi_i)$ , and  $\sigma_i = (s_i - s_{i0})$ , where  $s_{i0} \equiv \langle s_i \rangle_0$ . Then  $\mathcal{H}$  can be written as an infinite perturbation series in powers of the (perturbation) Hamiltonian part

$$S_f(s, \phi) = -\frac{1}{2} \sum_{ij} J_{ij} \sigma_i \sigma_j - \sum_{ij} \Delta_i \sigma_j. \quad (26)$$

The respective series can be presented in the form

$$\mathcal{H}_f(\phi) = -\frac{1}{2} \sum_{ij} J_{ij} \Delta_i \Delta_j + \sum_{l=1}^{\infty} \mathcal{H}_f^{(l)}(\phi), \quad (27)$$

where

$$\mathcal{H}_f^{(l)}(\phi) = \frac{(-\beta)^{l-1}}{l!} \langle S_f^l(s, \phi) \rangle_{0c}, \quad (28)$$

$\langle \dots \rangle_{0c}$  denotes the so-called connected averages [1]; for example, the average  $\langle S_f^2 \rangle_0 \langle S_f^2 \rangle_0$  is excluded from the connected  $\langle S_f^4 \rangle_{0c}$ . For this cumulant (semi-invariant) expansion rules, similar to the Wick theorem in the perturbation theory of propagator type, are not available and one should perform the calculations with some caution. The term  $\mathcal{H}_f^{(1)}$  is equal to zero, and this leads to a reduction of some infinite series in the next orders of the theory ( $l > 1$ ).

**2.2. Lowest Order Correction.** Let us consider the first term in the r.h.s. of Eq. (27) and neglect all others. This yields  $\mathcal{H} = (\mathcal{H}_0 + \mathcal{H}_f)$  in the form

$$\begin{aligned} \mathcal{H} \approx & \frac{\beta}{2} \sum_{ijk} J_{ij} J_{ik} \phi_j \phi_k - \frac{\beta^2}{2} \sum_{ijkl} J_{ij} J_{ik} J_{jl} \phi_k \phi_l - \\ & - \frac{\beta^3}{4} \sum_{ijklm} J_{ij} J_{ik} J_{il} J_{im} \phi_j \phi_k \phi_l \phi_m + \\ & + \frac{\beta^4}{3} \sum_{ijklmn} J_{ij} J_{jk} J_{il} J_{im} J_{in} \phi_k \phi_l \phi_m \phi_n. \end{aligned} \quad (29)$$

Performing this straightforward calculation one readily sees a very important property of the present theory, namely, that the first term in the r.h.s. of (15) is totally compensated by a respective counter term coming from the new contribution ( $\sim \Delta_i \Delta_j$ ) to the effective free energy. Besides, another new term *twice* compensates the second term in (15) so that the term of type  $JJ\phi\phi$  now appears with a positive sign. The  $\phi^4$ -part of the effective free energy also undergoes a drastic change due to the  $\Delta\Delta$ -correction coming from Eq. (28).

The same result can be obtained in a more general and, perhaps, more convenient way, if we add the  $\Delta\Delta$ -term in (27) to  $\mathcal{H}_0$  from (6) before doing the expansion of Landau type. Then, within the same lowest order approximation for series (27) we obtain a more general result for  $\mathcal{H}$ , namely

$$\begin{aligned} \mathcal{H} = & -\frac{1}{2} \sum_{ij} J_{ij} \text{th}(\beta a_i) \text{th}(\beta a_j) + \\ & + \sum_{ij} J_{ij} \phi_i \text{th}(\beta a_j) - \beta^{-1} \sum_i \ln [2\text{ch}(\beta a_i)]. \end{aligned} \quad (30)$$

This form of  $\mathcal{H}$  clearly shows the lack of the simple  $J_{ij} \phi_i \phi_j$  term describing the direct two-site exchange. In our further considerations we shall be faced only with inter-particle interactions (correlations), which are extended at distances larger than  $R_{\text{int}}$ . In order to obtain (29) one must expand the transcendental functions in (30);  $(\beta a_i) \ll 1$ .

New forms (29) and (30) of the effective free energy describe only indirect two-site interactions because the direct two-site interaction disappeared from our consideration. Now we are at a stage of description of correlations which extend up to  $2R_{ij}$  and larger distances. The tendency of the growing length scale of the interactions included in our effective free energy will be the main and unavoidable feature of our further consideration.

Another very important feature of our findings is that in the simplest variant of the theory when the field is uniform ( $\phi_i \equiv \phi$ ) as well as in the GL variant

in LWLA given by Eq. (24) the values of the Landau coefficients  $c_0$ ,  $r_0$ , and  $u_0$  do not change within the framework of accuracy of the effective  $\phi^4$ -field theory, where terms (Landau invariants) of order  $O(t^3)$  are ignored;  $t = (T - T_c)/T_c$ . This property seems to exist to any order of expansion (27) in the limit of an infinite-range ( $z \rightarrow \infty$ ) initial interaction  $J_{ij}$ . The  $1/z$ -corrections to the parameters ( $c_0, r_0, u_0$ ) of the effective field theory are obtained from the ( $l > 1$ )-terms in series (27).

**2.3. High-Energy Corrections of Higher Order.** We have already mentioned that the ( $l = 1$ )-term in (27) is zero. A calculation of the next two terms ( $l = 2, 3$ ) in (27) has been carried out in Ref. [2]. Following Ref. [2] we can write  $\mathcal{H}_f^{(2)}$  in the form

$$\mathcal{H}_f^{(2)} = -\frac{\beta}{4} \sum_{ij} \frac{J_{ij}^2}{\text{ch}^2(\beta a_i) \text{ch}^2(\beta a_j)} - \frac{\beta}{2} \sum_{ijk} J_{ij} J_{ik} \frac{\Delta_j \Delta_k}{\text{ch}^2(\beta a_i)}. \quad (31)$$

Result (31) gives the first ( $1/z$ )-corrections to the parameters  $c_0, r_0$ , and  $u_0$ . Adding result (31) to  $\mathcal{H}(\phi)$  we obtain a form of the effective Hamiltonian  $\mathcal{H}(\phi)$  which is more precise than the preceding ones. Let us write down this quite lengthy expression of the effective Hamiltonian:

$$\begin{aligned} \mathcal{H} = & \frac{\beta^2}{2} \sum_{ijkl} J_{ij} J_{ik} J_{jl} \phi_k \phi_l + \frac{\beta^3}{2} \sum_{ijkl} J_{ij}^2 J_{ik} J_{il} \phi_k \phi_l - \\ & - \frac{\beta^3}{2} \sum_{ijklm} J_{ij} J_{ik} J_{jl} J_{km} \phi_l \phi_m + \frac{\beta^3}{4} \sum_{ijklm} J_{ij} J_{ik} J_{il} J_{im} \phi_j \phi_k \phi_l \phi_m - \\ & - \beta^4 \sum_{injkml} J_{in} J_{ij} J_{ik} J_{il} J_{nm} \phi_j \phi_k \phi_l \phi_m + \\ & + \frac{\beta^5}{3} \sum_{inpjklm} J_{in} J_{ip} J_{nj} J_{nk} J_{nl} J_{pm} \phi_j \phi_k \phi_l \phi_m + \\ & + \frac{\beta^5}{2} \sum_{inpjklm} J_{in} J_{ip} J_{ij} J_{ik} J_{nl} J_{pm} \phi_j \phi_k \phi_l \phi_m - \\ & - \frac{\beta^5}{3} \sum_{injkml} J_{in}^2 J_{ij} J_{ik} J_{il} J_{im} \phi_j \phi_k \phi_l \phi_m - \\ & - \frac{\beta^5}{4} \sum_{injkml} J_{in}^2 J_{ij} J_{ik} J_{il} J_{im} \phi_j \phi_k \phi_l \phi_m. \quad (32) \end{aligned}$$

Result (32) shows that the indirect two-point interactions of type  $JJ\phi\phi$  available in effective Hamiltonian (29) do not exist in this higher accuracy of the theory.

The interactions of type  $JJJ\phi\phi$  and  $JJJJ\phi\phi$  presented in effective Hamiltonian (32) extend up to distances  $3R_{\text{int}}$ . The same is valid for the four point interactions included in (32).

Within LWLA Eq. (32) yields

$$\mathcal{H} = \frac{1}{2} \sum_{\mathbf{k}} (r + ck^2) |\phi(\mathbf{k})|^2 + \frac{u}{N} \sum_{(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)} \phi(\mathbf{k}_1)\phi(\mathbf{k}_2)\phi(\mathbf{k}_3)\phi(-\mathbf{k}_1 - \mathbf{k}_2 - \mathbf{k}_3). \quad (33)$$

In (33),

$$c = \left(1 + \frac{5}{z}\right) c_0, \quad r = \left(1 + \frac{3}{z}\right) \tilde{r}_0, \quad u = \left(1 + \frac{4}{z}\right) u_0, \quad (34)$$

where  $\tilde{r}_0 = k_B(T - T_c)$  is given by the «true» (renormalized) critical temperature

$$T_c = T_{c_0} \left(1 - \frac{1}{z}\right). \quad (35)$$

In deriving this lattice version of the effective Hamiltonian we have performed the lattice summations in (32) in the reciprocal ( $\mathbf{k}$ )-space with the help of LWLA:  $J(k) \approx (J - c_0k^2)$ .

The present results demonstrate a type of renormalization of the GL parameters ( $c_0, r_0, u_0$ ) of the effective Hamiltonian due to  $1/z$ -corrections. By a suitable choice of units, one of these parameters can be kept invariant, for example, equal to unity. Therefore, within a suitable normalization of the theory, the field  $\phi(\mathbf{x})$  acquires a  $1/z$ -correction as well [2].

The term  $\mathcal{H}_f^{(3)}$  in (27) has the form

$$\begin{aligned} \mathcal{H}_f^{(3)} = & -\frac{\beta^2}{3} \sum_{ij} J_{ij}^3 \frac{\text{th}(\beta a_i) \text{th}(\beta a_j)}{\text{ch}^2(\beta a_i) \text{ch}^2(\beta a_j)} \\ & - \frac{\beta^2}{6} \sum_{ijl} \frac{J_{ij} J_{il} J_{jl}}{\text{ch}^2(\beta a_i) \text{ch}^2(\beta a_j) \text{ch}^2(\beta a_l)} \\ & - \beta^2 \sum_{ijl} J_{ij}^2 J_{jl} \frac{\Delta_l \text{th}(\beta a_j)}{\text{ch}^2(\beta a_i) \text{ch}^2(\beta a_j)} \\ & - \frac{\beta^2}{2} \sum_{ijln} J_{ij} J_{il} J_{jn} \frac{\Delta_l \Delta_n}{\text{ch}^2(\beta a_i) \text{ch}^2(\beta a_j)} \\ & - \frac{\beta^2}{3} \sum_{ijln} J_{ij} J_{il} J_{in} \frac{\Delta_j \Delta_l \Delta_n \text{th}(\beta a_i)}{\text{ch}^2(\beta a_i)}. \end{aligned} \quad (36)$$

Let us consider the contribution of the term  $\mathcal{H}_f^{(3)}$  to the quadratic ( $\phi^2$ ) part of effective Hamiltonian (33). In performing the calculations for both short-range ( $nn$ ) and long-range ( $R_{\text{int}} \gg a_0$ ) we have to evaluate again several lattice sums. Here we shall mention a particular sum, namely,

$$\frac{1}{N} \sum_{ijl} J_{ij} J_{il} J_{jl}, \quad (37)$$

which is equal to zero for  $nn$  interactions but gives a contribution ( $\approx J^3/z$ ) for interaction radius  $R_{\text{int}} > a_0$ . Thus we introduce the following interpolation formula:  $E = \kappa(z)/z$ , where  $0 \leq \kappa(z) \leq 1$  is an interpolation parameter which is supposed to be a smooth function of the coordination number  $z$ . The limiting case  $\kappa = 0$  corresponds to  $nn$  interactions and the limiting case  $\kappa = 1$  corresponds to interactions of larger size. We suppose that the shape of the function  $\kappa(z)$  depends on details of the function  $J(R)$ .

Bearing in mind these notes we have calculated the quadratic ( $\phi_i \phi_j$ ) contribution to the effective Hamiltonian  $\mathcal{H}(\phi)$  which comes from the Hamiltonian part  $\mathcal{H}_f^{(3)}(\phi)$  given by (36). Here we present the results for the parameters  $T_{c_0}$ ,  $c$ , and  $r$  which define the quadratic part  $\mathcal{H}_2$  of  $\mathcal{H}$ :

$$\mathcal{H}_2(\phi) = \frac{1}{2} \sum_{\mathbf{k}} G_0^{-1}(k) |\phi(\mathbf{k})|^2, \quad (38)$$

with the (bare) correlation function

$$G_0^{-1}(k) = \mathcal{D}_1(z) [T - T_c + \mathcal{D}_2(z) T_c \rho^2 k^2]. \quad (39)$$

Here

$$T_c = \left( 1 - \frac{1 + \kappa}{z} - \frac{1 + 3\kappa}{3z^2} \right) T_{c_0}, \quad (40)$$

$$\mathcal{D}_1(z) = 1 + \frac{3 + 4\kappa}{z} + \frac{22 + 60\kappa + 30\kappa^2}{3z^2}, \quad (41)$$

and

$$\mathcal{D}_2(z) = 1 + \frac{2 + 3\kappa}{z} + \frac{14 + 30\kappa + 9\kappa^2}{3z^2}. \quad (42)$$

Result (35) for  $T_c(z)$  has been published for the first time [3] in case of  $nn$  interactions ( $\kappa = 0$ ); note, that there are errors in Ref. [3] for the functions  $\mathcal{D}_1(z)$  and  $\mathcal{D}_2(z)$ .

The functions  $\mathcal{D}_1(z)$  and  $\mathcal{D}_2(z)$  renormalize the field  $\phi(\mathbf{k})$  and the vertices  $c_0$ , and  $r_0$ . For a total renormalization of the parameters of the theory up to the second order in the  $1/z$ -expansion we need to know the  $(1/z)^2$ -correction to the vertex  $u_0$ . We suppose that the calculation of this correction can be accomplished

in the way described above; this « $z$ -renormalization» has been discussed to the first order in  $(1/z)$  in Ref. [2]. Here we wish to stress that within our extension of the theory the magnetic susceptibility  $G_0(0)$  is  $\mathcal{D}_1$  times smaller than the known MF susceptibility corresponding to  $\mathcal{D}_1(\infty) = 1$ .

The numerical coefficients in (40)–(42) indicate that real numbers  $z$  of  $nn$  like  $n = 2, 4, 6$  for simple lattices of spatial dimensionalities  $D = 1, 2, 3$ , respectively, give a good expansion parameter  $1/z$ . The  $1/z$ -corrections are more substantial for the case of short-range interactions ( $R_{\text{int}} \sim a_0$ ), and one may suppose that for such interactions the  $(1/z)$ -series (40)–(42) are asymptotic; for the case of  $T_c$ , see a discussion of this topic in Ref. [8]. But even in case of asymptotic type of these series they may give more reliable results than the «bare» values  $(c_0, r_0, u_0)$  of the Landau parameters; see arguments presented in Ref. [8].

**2.4. Critical Temperature.** The critical temperature  $T_c$  given by (40) can be compared with exact and reliable numerical (MC) results. Let us consider  $nn$  interactions ( $\kappa = 0$ ). For one-dimensional (1D) IM, we know that  $T_c = 0$ , MFA predicts  $T_c = 2J_0/k_B$  (in this case,  $z = 2D = 2$ ), and (40) yields  $T_c = 5J_0/6k_B$ . This is a quite good result for 1D systems with very strong fluctuation effects. In 2D systems the fluctuations are not so strong and we find that (40) reproduces the exact Onsager result ( $T_c = 2.27J_0/k_B$ ) with an error of 22%, i.e., we have  $T_c = 35J_0/12k_B$ . For 3D systems our result is  $T_c = 89J_0/18k_B$ , whereas the best series analysis and MC results yield a difference of 9%:  $T_c = 4.5J_0/k_B$  (see, e.g., Refs. [16, 17]). Our results seem quite reliable.

**2.5. Ground State.** The  $1/z^2$ -correction to the vertex  $u_0$  has not yet been calculated although this calculation does not present difficulties. In this situation we shall give notion for the ground state energy by using the first order corrections to  $r_0$  and  $u_0$ . The equilibrium free energy per site  $f = (F/N)$  is given by  $F = \mathcal{H}$  and (33) for the ( $k = 0$ )-Fourier amplitude  $\phi(0)$ , which minimizes  $f$ . Denote for convenience  $\phi(0) = \sqrt{N}\varphi \neq 0$  for the low-temperature ordered phase. From (33) we obtain  $f = -(r^2/16u) < 0$  whereas the «unrenormalized» free energy is  $f_0 = (\mathcal{H}_0/N) = -(r_0^2/16u_0) < 0$ . Thus, using (34) we have  $f(T) = (1 + 2/z)f_0(T)$ , which means that the new effective theory has a lower energy of the ordered phase. This is true also in the case of  $T = 0$ , where  $r_0 = -k_B T_{c_0} = -J$ . For the zero temperature (ground) state we have  $f_0(0) = -3J/4$  and  $f(0) = -3(1 + 2/z)J/4$ . This result is also along the right direction because the MF theories ( $\mathcal{H}_0$ ) give unreliable high values of the ground state energy. The order parameter  $\varphi^2(T) = (-r/4u)$  is  $(1 - z)$  times smaller than the respective quantity  $\varphi_0^2(T) = (-r_0/4u_0)$  in the usual theory based on  $\mathcal{H}_0$ .

**2.6. Effective Interactions and Phenomenon of Growing of Statistical Correlations.** We have explicitly shown the phenomenon of the growing length size of the interparticle correlations in a classic system of interacting particles. To see this we have already introduced a new interpretation of the terms in the effective Hamiltonian (see Subsec. 1.4). Let us consider the terms present in  $\mathcal{H}$  as



terms describing concrete intersite interactions. While the initial interaction  $J_{ij}$  in IM ensures only two-site correlations (interactions), effective Hamiltonians (15), (29), and (32) contain multi-site effective interactions. In contrast to usual theory (15), where only extremely short-range effective correlations are contained, the more precise effective Hamiltonians contain long-range two-site ( $\phi_i\phi_j$ ) and four-site ( $\phi_i\phi_j\phi_k\phi_l$ ) correlations, and all of these correlations are indirect, i.e. the correlation, for example, between two sites ( $ij$ ) is mediated by one or more other sites ( $k, \dots$ ). A direct ( $J_{ij}$ )-interaction is presented by the first term in the r.h.s. of (15) but also the system exhibits two indirect correlations of type  $\phi_i\phi_j$  and  $\phi_i\phi_j\phi_k\phi_l$  given by the last two terms in the r.h.s. of (15). In the more precise variants of the theory, where a larger portion of the initial partition function has been calculated, the direct intersite interaction vanishes, and the particles are correlated only by indirect effective interactions. The length scale of these correlations grows in a monotonous way with the increase of the accuracy of the calculation, i.e. with the increase of the number  $l$  of the terms in series (27). If we take the two-site correlations in the  $nn$  IM as an example, the maximal length of extension of these correlations in (1) is  $a_0$ , in (15) –  $2a_0$ , in (29) –  $3a_0$ , in (32) –  $4a_0$ , i.e.  $(p-1)a_0$ , where  $p$  is the maximal number of the summation indices in the terms of type  $\phi_i\phi_j$  in a given effective Hamiltonian. Surely, the number  $p$  tends to  $N$ . This means that the most accurate effective field theory of many-body systems will correspond to (almost-) infinite range of correlations.

The origin of these correlations is purely statistical. This effect is known and has both general formulation and application in many-body physics. Here we have established and described in detail the concrete mechanism of this effect and, moreover, we have performed a demonstration of the remarkable picture of successive growth of the correlation length scale.

**2.7. Final Remarks.** Obviously the  $(1/z)$ -corrections are not the main point of discussion at the end of this paper of restricted length. Let us mention that the growth of the correlations discussed in Subsec.2.6 is not related with the  $1/z$ -corrections. It exists for any  $z$ , even in the «MF limiting case» of  $z \rightarrow \infty$ , when the GL parameters keep their «initial» values  $(c_0, r_0, u_0)$ . This effect follows from the fact that the terms in the initial Hamiltonian are compensated by the first «fluctuation» correction; see the first term in the r.h.s. of (27). At the next level of accuracy of the calculation, terms coming from the  $(l=2)$ -term in (27) compensate the available terms and this process continues up to the incorporation of all particles in the correlation phenomenon; remember that the term corresponding to  $l=1$  is equal to zero.

Here we emphasize that the terms in  $\mathcal{H}_0$  – actually one of the most often used Hamiltonian, do not exist at all. They vanish just after the inclusion of the first correction to the usual theory; see  $\Delta_i\Delta_j$ -correction in (27). In place of these terms other terms with more complex structure come from perturbation series (27). The outlined picture clearly indicates, that the terms which finally remain

in the  $\phi^4$ -theory, are terms of type

$$\frac{1}{2} (\beta^{M-1} J^M - \beta^M J^{M+1}) \phi^2, \quad M \sim N, \quad (43)$$

where obvious notations have been introduced; for example,  $\beta^1 J^2$  denotes the first term in the r.h.s. of (29). The  $\phi^4$ -terms behave differently, because the lowest order term in  $J$ , namely, a term of type  $\beta^3 J^4 \phi^4$  appears at any step of development of series (27).

At any stage of this surprising picture of the infinite series of successive modifications of the Hamiltonian both  $\phi^2$ - and  $\phi^4$ -terms keep their numerical coefficients equal to that in the usual GL Hamiltonian  $\mathcal{H}_0$ . This is true within the whole scope of validity of the expansion in powers of  $\phi$ .

Our consideration justifies the GL fluctuation Hamiltonian. However, we have presented a new and quite surprising picture of the interparticle correlations, which reveals new remarkable properties of the GL theory. Apart from the  $1/z$ -corrections to the GL parameters, the structure of this theory is absolutely comprehensive as a tool for investigation of large-scale correlation phenomena in many-body systems. We are certain that our findings have an application beyond the field of phase transitions.

**Acknowledgments.** The author thanks Professor N.M. Plakida and the staff of the Bogoliubov Laboratory of Theoretical Physics (JINR) for the hospitality during a short-term visit. Financial support by a research grant (JINR) and Scenet-EC (Parma) is also acknowledged.

## REFERENCES

1. *Uzunov D. I.* Theory of critical phenomena. Singapore: World Scientific, 1993.
2. *Uzunov D. I.* Lectures on cooperative phenomena in condensed matter. Sofia: Heron Press, 1996. P. 46–110.
3. *Uzunov D. I.* Coherent approaches to fluctuations. Singapore: World Scientific, 1996. P. 25.
4. *Brout R.* // Phys. Rev. 1959. V. 115. P. 824.
5. *Horwitz G., Callen H. B.* // Phys. Rev. 1961. V. 124. P. 1757.
6. *Englert F.* // Phys. Rev. 1963. V. 129. P. 567.
7. *Stinchcombe R. B., Horwitz G., Englert F., Brout R.* // Phys. Rev. 1963. V. 130. P. 155.
8. *Fisher M. E., Gaunt D. S.* // Phys. Rev. 1964. V. 113. P. A224.
9. *Brout R.* Phase transitions. N.Y.; Amsterdam: University of Brussels, 1965.

10. *Vaks V. G., Larkin A., Pikin S. A.* // J. Exp. Teor. Fiz. 1966. V.51. P.361 [Russ. Phys. — JETP. 1967. V.24. P.240].
11. *Vaks V. G., Larkin A. I., Pikin S. A.* // J. Exp. Teor. Fiz. 1967. V.53. P.281.
12. *Vaks V. G., Larkin A. I., Pikin S. A.* // J. Exp. Teor. Fiz. 1967. V.53. P.1089.
13. *Brout R.* // Phys. Rep. C. 1974. V.10. P.1.
14. *Kirkwood J. G.* // J. Chem. Phys. 1938. V.6. P.70.
15. *Uzunov D. I.* In preparation. 2004.
16. *Baker G. A., Jr.* // J. Stat. Phys. 1994. V.77. P.955.
17. *Baker G. A., Jr.* Coherent approaches to fluctuations. Singapore: World Scientific, 1996. P.19.

Received on November 15, 2004.

Корректор *Т. Е. Попеко*

Подписано в печать 09.03.2005.

Формат 60 × 90/16. Бумага офсетная. Печать офсетная.

Усл. печ. л. 1,43. Уч.-изд. л. 2,04. Тираж 305 экз. Заказ № 54814.

Издательский отдел Объединенного института ядерных исследований  
141980, г. Дубна, Московская обл., ул. Жолио-Кюри, 6.

E-mail: [publish@pds.jinr.ru](mailto:publish@pds.jinr.ru)

[www.jinr.ru/publish/](http://www.jinr.ru/publish/)