

E1-2007-166

L. S. Azhgirey, A. V. Tarasov

CALCULATION OF SPIN ALIGNMENT OF DEUTERONS
TRAVELING THROUGH MATTER

Submitted to «Particles and Nuclei, Letters»

Ажгирей Л. С., Тарасов А. В.

E1-2007-166

Вычисление выстроенности проходящего через вещество пучка дейтронов

Описан расчет выстроенности по спине пучка релятивистских дейтронов, появляющейся при его прохождении через вещество. Тензорная поляризация вычисляется в рамках теории многократного рассеяния Глаубера. Результаты вычислений сравниваются с полученными недавно экспериментальными данными.

Работа выполнена в Лаборатории ядерных проблем им. В. П. Дзелепова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2007

Azhgirey L. S., Tarasov A. V.

E1-2007-166

Calculation of Spin Alignment of Deuterons Traveling through Matter

Calculations of the spin alignment of the relativistic deuteron beam passing through matter are described. The tensor polarization is calculated within the framework of the Glauber multiple scattering theory. The calculation results are compared with the recent experimental data.

The investigation has been performed at the Dzhelepov Laboratory of Nuclear Problems, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 2007

1. It is known that the availability of a small quadruple moment of the deuteron gives rise to a number of polarization effects in the nuclear reactions involving the deuteron [1–5]. Recently the phenomenon of spin dichroism (production of a spin-aligned deuteron beam arising as initially unpolarized deuterons pass through matter) has been observed using an extracted unpolarized 5-GeV/ c deuteron beam of the Nuclotron [6]. In this report the results of the calculation of the tensor polarization of the relativistic deuterons traveling through matter are given.

2. Before writing the expression for the total cross section of the deuteron-nucleus scattering, let us consider the nucleon-deuteron (ND) scattering. The total cross section of the ND scattering for a definite spin state of the deuteron λ may be represented in the form [7, 8]

$$\sigma_{ND}^{\text{tot}(\lambda)} = 2\text{Re} \int d^3r_D \psi_D^{(\lambda)\dagger}(\mathbf{r}_D) \psi_D^{(\lambda)}(\mathbf{r}_D) \int d^2b \gamma_{ND}(\mathbf{b}, \mathbf{s}). \quad (1)$$

Here $\mathbf{r}_D = (\mathbf{s}_D, z_D)$ is the distance between the nucleon and deuteron center of gravity, \mathbf{s}_D and z_D are the transversal and longitudinal parts of this distance, so that $d^3r_D = d^2s_D dz_D$.

The profile function $\gamma_{ND}(\mathbf{b}, \mathbf{s})$ can be expressed through the functions γ_{Np} and γ_{Nn} for the proton and the neutron, respectively:

$$\begin{aligned} \gamma_{ND}(\mathbf{b}, \mathbf{s}_D) = & \left[\gamma_{Np} \left(\mathbf{b} - \frac{1}{2} \mathbf{s}_D \right) + \gamma_{Nn} \left(\mathbf{b} + \frac{1}{2} \mathbf{s}_D \right) - \right. \\ & \left. - \gamma_{Np} \left(\mathbf{b} - \frac{1}{2} \mathbf{s}_D \right) \gamma_{Nn} \left(\mathbf{b} + \frac{1}{2} \mathbf{s}_D \right) \right]. \quad (2) \end{aligned}$$

The nucleon profile functions are connected with the corresponding amplitudes of NN scattering by the expressions

$$\begin{aligned} \gamma_{Np} \left(\mathbf{b} - \frac{1}{2} \mathbf{s}_D \right) &= \int d^2q f_{Np}(\mathbf{q}) \exp \left[-i \left(\mathbf{b} - \frac{1}{2} \mathbf{s}_D \right) \mathbf{q} \right], \\ \gamma_{Nn} \left(\mathbf{b} + \frac{1}{2} \mathbf{s}_D \right) &= \int d^2q f_{Nn}(\mathbf{q}) \exp \left[-i \left(\mathbf{b} + \frac{1}{2} \mathbf{s}_D \right) \mathbf{q} \right]. \end{aligned} \quad (3)$$

If the amplitudes of the NN scattering at high energies have the form

$$\begin{aligned} f_{Np}(\mathbf{q}) &= \frac{i}{4\pi} \sigma_{Np}^{\text{tot}} (1 - i\alpha_{Np}) \exp \left[-\frac{1}{2} Bq^2 \right], \\ f_{Nn}(\mathbf{q}) &= \frac{i}{4\pi} \sigma_{Nn}^{\text{tot}} (1 - i\alpha_{Nn}) \exp \left[-\frac{1}{2} Bq^2 \right], \end{aligned} \quad (4)$$

the profile functions may be written in the form

$$\begin{aligned} \gamma_{Np} \left(\mathbf{b} - \frac{1}{2} \mathbf{s}_D \right) &= \frac{1}{4\pi B} (1 - i\alpha_{Np}) \exp \left[-\frac{1}{2B} \left(\mathbf{b} - \frac{1}{2} \mathbf{s}_D \right)^2 \right], \\ \gamma_{Nn} \left(\mathbf{b} + \frac{1}{2} \mathbf{s}_D \right) &= \frac{1}{4\pi B} (1 - i\alpha_{Nn}) \exp \left[-\frac{1}{2B} \left(\mathbf{b} + \frac{1}{2} \mathbf{s}_D \right)^2 \right], \end{aligned} \quad (5)$$

The total cross section of NN scattering σ_{NN} and the ratio of the real to imaginary parts of the forward NN scattering amplitude α averaged over the deuteron nucleons are

$$\sigma_{NN} = \frac{1}{2} (\sigma_{pp} + \sigma_{pn}), \quad \alpha = \frac{\sigma_{pp} \alpha_{pp} + \sigma_{pn} \alpha_{pn}}{\sigma_{pp} + \sigma_{pn}}. \quad (6)$$

The total cross section of the deuteron scattering on a nucleus is

$$\begin{aligned} \sigma_{DA}^{\text{tot}(\lambda)} &= 2\text{Re} \int d^3 r_D \psi_D^{(\lambda)\dagger}(\mathbf{r}_D) \psi_D^{(\lambda)}(\mathbf{r}_D) \int d^2 b \times \\ &\quad \times \int |\Psi_A(\{r_i\})|^2 \prod_{i=1}^A d^3 r_i \Gamma_{AD}(\mathbf{b}, \mathbf{s}_D; \{\mathbf{s}_i\}). \end{aligned} \quad (7)$$

The nuclear profile function has the form

$$\Gamma_{AD} = 1 - \prod_{i=1}^A \left[1 - \gamma_{ND}(\mathbf{b} - \mathbf{s}_i, \mathbf{s}_D) \right]. \quad (8)$$

Here \mathbf{b} is the impact parameter between the deuteron and the nucleus, \mathbf{s}_i and \mathbf{s}_D are the transversal parts of the distances of the nucleus nucleon from the center of gravity of the nucleus and the deuteron nucleon from the center of gravity of the deuteron, respectively.

The wave function of a nucleus $\Psi_A(\{\mathbf{r}_i\})$ is connected with the nuclear density $\rho(\mathbf{r}_i)$ by the relation

$$|\Psi_A(\{\mathbf{r}_i\})|^2 = \prod_{i=1}^A \rho(\mathbf{r}_i), \quad (9)$$

where

$$\int \rho(\mathbf{r}) d^3 r = 1, \quad \rho(\mathbf{r}) = \rho(\mathbf{s}, z). \quad (10)$$

The second integral in the expression for the total DA cross section may be written in the form

$$\begin{aligned} \int |\Psi_A(\{r_i\})|^2 \prod_{i=1}^A d^3 r_i \Gamma_{AD}(\mathbf{b}, \mathbf{s}_D; \{\mathbf{s}_i\}) &= \\ &= 1 - \left[1 - \int \rho(\mathbf{s}, z) d^2 s dz \Gamma_{AD}(\mathbf{b}, \mathbf{s}_D; \mathbf{s}) \right]^A. \end{aligned} \quad (11)$$

The spin structure of the deuteron wave function is expressed in the following way:

$$\psi_D^{(\lambda)}(\mathbf{r}) = \left[\phi_s(\mathbf{r}) + \frac{1}{2\sqrt{2}} (3 \sigma_n \nu \cdot \sigma_p \nu - 1) \phi_d(\mathbf{r}) \right] \chi_D^{(\lambda)}, \quad (12)$$

where

$$\nu = \frac{\mathbf{r}}{r}, \quad \int \{ |\phi_s(\mathbf{r})|^2 + |\phi_d(\mathbf{r})|^2 \} d^3 r = 1.$$

Here $\phi_s(\mathbf{r})$ and $\phi_d(\mathbf{r})$ are the wave functions of the S - and D -states of the deuteron, ν is the orientation of the deuteron spin. The spin function of the deuteron $\chi_D^{(\lambda)}$ is connected with the nucleon spin functions by the relations

$$\begin{aligned} \chi_D^{(+1)} &= \chi_p^{(1/2)} \chi_n^{(1/2)}, \\ \chi_D^{(-1)} &= \chi_p^{-(1/2)} \chi_n^{-(1/2)}, \\ \chi_D^{(0)} &= \frac{1}{\sqrt{2}} \left[\chi_p^{(1/2)} \chi_n^{(-1/2)} + \chi_p^{(-1/2)} \chi_n^{(1/2)} \right]. \end{aligned} \quad (13)$$

3. If the deuteron wave functions ϕ_s and ϕ_d can be represented as the sums of the Gauss functions [10]

$$\phi_s = \sum_{i=1}^5 A_i \exp(-\alpha_i q^2), \quad \phi_d = q^2 \sum_{i=1}^5 B_i \exp(-\beta_i q^2), \quad (14)$$

and the nuclear density is chosen in the simple Gaussian form

$$\rho(\mathbf{r}) = \frac{1}{(\pi R_0^2)^{3/2}} \exp\left(-\frac{r^2}{R_0^2}\right), \quad (15)$$

the integrals of this problem can be taken analytically.

4. On the above assumptions in line with the multiple scattering theory, the difference of the total cross sections for the scattering of deuterons in different spin states (± 1) and (0) from the nuclei may be written in the form

$$\Delta\sigma = \frac{1}{N_S + N_D} \sum_{N=1}^A (-1)^{N-1} \frac{A!}{(A-N)!} \Delta\sigma^{(N)}, \quad (16)$$

where the cross section difference for the N th collision is given by

$$\begin{aligned} \Delta\sigma^{(N)} &= \pi R_1 R_2 \sum_{m=0}^N \sum_{n=0}^{N-m} \times \\ &\times \frac{\Delta_{m,n}^{(N)} a_1^{m+n} a_2^{N-m-n} \Omega_{m,n}^{(N)}}{[(m+n)R_2 + (N-m-n)R_1] n! m! (N-m-n)!}. \end{aligned} \quad (17)$$

There the following notations were used:

$$\begin{aligned} \Delta_{m,n}^{(N)} &= 3 \sum_{i=1}^5 \sum_{k=1}^5 C_i D_k \left(\frac{\pi}{\tau_{i,k}} \right)^{3/2} \frac{\lambda_{m,n}^{(N)}}{(\lambda_{m,n}^{(N)} + \tau_{i,k})^2} + \\ &+ \frac{3}{2} \sum_{i=1}^5 \sum_{k=1}^5 D_i D_k \left(\frac{\pi}{\nu_{i,k}} \right)^{3/2} \frac{\lambda_{m,n}^{(N)} (3\lambda_{m,n}^{(N)} + 7\nu_{i,k})}{\nu_{i,k} (\lambda_{m,n}^{(N)} + \nu_{i,k})^3} \end{aligned}$$

with

$$\begin{aligned} \lambda_{m,n}^{(N)} &= \frac{1}{4} \left(\frac{N-m-n}{B} + \frac{4mnR_2^2 + (m+n)(N-m-n)R_1^2}{R_1 [(m+n)R_2^2 + (N-m-n)R_1^2]} \right), \\ \Omega_{m,n}^{(N)} &= \cos(2N-m-n)\Phi, \quad \Phi = \arctan \alpha. \end{aligned} \quad (18)$$

The parameters R_1 , R_2 , a_1 and a_2 are expressed in terms of constants peculiar to this problem:

$$\begin{aligned} R_1^2 &= R_0^2 + 2B, \quad R_2^2 = R_0^2 + B, \\ a_1 &= \frac{\sigma_{NN}}{2\pi R_1^2} \sqrt{1 + \alpha^2}, \quad a_2 = -\frac{\sigma_{NN}^2 (1 + \alpha^2)}{16\pi^2 B R_2^2}, \\ R_0^2 &= \frac{2}{3} [\langle r_A^2 \rangle - \langle r_p^2 \rangle], \end{aligned} \quad (19)$$

where $\langle r_A^2 \rangle$ and $\langle r_p^2 \rangle$ are the squares of the rms radii of the nucleus and the proton, respectively.

The effective numbers for the S - and D -states are

$$N_S = \sum_{i=1}^5 \sum_{k=1}^5 \frac{C_i C_k \pi^{3/2}}{(\rho_i + \rho_k)^{3/2}}, \quad N_D = \frac{15}{8} \sum_{i=1}^5 \sum_{k=1}^5 \frac{D_i D_k \pi^{3/2}}{(\omega_i + \omega_k)^{7/2}}, \quad (20)$$

where

$$\begin{aligned}
C_i &= A_i (2.5/\alpha_i)^{3/2}, & D_i &= \sqrt{2}B_i (2.5/\beta_i)^{7/2}, \\
\rho_i &= 6.25/\alpha_i, & \omega_i &= 6.25/\beta_i, \\
\tau_{i,k} &= \rho_i + \omega_k, & \nu_{i,k} &= \omega_i + \omega_k.
\end{aligned}$$

5. The following values of the parameters were used in the calculations: $\sigma_{NN} = 4.40 \text{ fm}^2$, $\alpha_{NN} = -0.339$, $B = 0.297 \text{ fm}^2$ [9]. The constants a_i , b_i , α_i and β_i were taken from [10]; they correspond to the deuteron wave function of the Reid soft core potential. The rms radii used in the calculations are given in the second column of the table [11].

Root-mean-square radii R_A , cross section differences $\Delta\sigma$, P_{ZZ} , and target thicknesses x for different nuclei A calculated under the assumption that the beam intensity behind the target falls down to 0.01 of the initial one


A	R_A , fm	σ_t , mb	$\Delta\sigma$, fm ²	P_{ZZ}	x , cm
⁹ Be	2.26	540	3.16	0.171	68.8
¹² C	2.42	650	3.79	0.174	83.0
²⁷ Al	3.06	1090	6.25	0.170	70.3
⁴⁰ Ca	3.52	1400	7.78	0.165	141.2
⁶⁴ Cu	3.88	1880	9.58	0.151	29.0
¹⁰⁶ Ag	4.40	2630	11.91	0.135	30.0
¹⁹⁷ Au	5.33	3850	15.56	0.121	20.3
²⁰⁷ Pb	5.42	3980	15.93	0.120	35.2

6. It can be shown that the tensor polarization of the deuteron beam arising from this cross section difference is

$$P_{ZZ} = \frac{1 - \exp(-N\Delta\sigma x)}{1 + \frac{1}{2} \exp(-N\Delta\sigma x)}, \quad (21)$$

where N is the number of nuclei in cm³ of matter x cm thick. The calculation results are shown in the figure by solid curve together with the recent experimental data [6].

In the table the values of P_{ZZ} for different nuclei were calculated under the assumption that the beam intensity behind the target falls down to 0.01 of the initial one. The total cross sections of the deuteron nucleus interaction σ_t were calculated according to [12].



Tensor polarization of deuterons vs thickness of the carbon target [6]. The dashed region shows the error corridor, the solid curve is the calculation result

7. A formalism is elaborated to calculate the tensor polarization of an initially unpolarized deuteron beam arising as deuterons pass through matter. The effect is treated within the framework of the Glauber multiple scattering theory.

The results of the calculation are compared with the recent experimental data obtained with the extracted unpolarized 5-GeV/ c deuteron beam of the Nuclotron. The calculation results are in qualitative agreement with the experimental data.

The possibility of producing spin-aligned high-energy deuteron beams at the sacrifice of beam intensity is noted.

Acknowledgments. The authors express their gratitude to Professors V. G. Baryshevsky and L. S. Zolin for helpful discussions. The research described in this publication was supported in part by the Russian Foundation for Basic Research under grant No. 06-02-16728.

REFERENCES

1. Franco V., Glauber R. J. // Phys. Rev. Lett. 1969. V. 22. P. 370.
2. Harrington D. // Phys. Lett. B. 1969. V. 29. P. 188.
3. Azhgirey L. S. et al. // Phys. Lett. B. 1995. V. 361. P. 21.
4. Baryshevsky V. G. // J. Phys. G: Nucl. Part. Phys. 1993. V. 19. P. 273.
5. Baryshevsky V. et al. arXiv:hep-ex/0501045 v2 (2005).
6. Azhgirey L. S. et al. JINR Preprint E1-2007-165. Dubna, 2007 (in press).

7. *Glauber R. J.* // Lectures in Theoretical Physics / Ed. by W. E. Brittin et al. V. 1. New York: Interscience Publishers, Inc., 1959. P. 315.
8. *Franco V., Glauber R. J.* // Phys. Rev. 1966. V. 142. P. 1195.
9. pdg.lbl.gov/2006/tables.html
10. *Alberi G., Rosa L. P., Thomé Z. D.* // Phys. Rev. Lett. 1975. V. 34. P. 503.
11. *Elton L. R. B.* Nuclear Sizes. Oxford University Press, 1961.
12. *Fäldt G., Pilkuhn H.* // Ann. Phys. 1970. V. 58. P. 454.

Received on November 1, 2007.

Редактор *Е. И. Кравченко*

Подписано в печать 26.12.2007.

Формат 60 × 90/16. Бумага офсетная. Печать офсетная.

Усл. печ. л. 0,62. Уч.-изд. л. 0,9. Тираж 365 экз. Заказ № 56010.

Издательский отдел Объединенного института ядерных исследований
141980, г. Дубна, Московская обл., ул. Жолио-Кюри, 6.

E-mail: publish@jinr.ru

www.jinr.ru/publish/